

University of California IPFP, IPM, and WPMA Pest Update #6
Tehama County 4/23/07

IPFP = Integrated Prune Farming Practices
 IPM = Integrated Pest Management
 WPMA = Walnut Pest Management Alliance

| INSECT | FIRST BIOFIX | SECOND BIOFIX | THIRD BIOFIX | FOURTH BIOFIX | ACTIVITY MOTHS/DAY | AVG. DEG. DAYS/DAY | DAY DEGREES FROM BIOFIX |
|---------------------|--------------|---------------|--------------|---------------|--------------------|--------------------|-------------------------|
| CODLING MOTH | 3/26 | - | - | - | 0 | 3.5 | 200 |
| ORIENTAL FRUIT MOTH | 3/5 | - | - | - | 2.9 | 6.8 | 613 |
| PEACH TWIG BORER | 4/2 | - | - | - | 1.9 | 3.3 | 152 |
| NAVEL ORANGE WORM | - | - | - | - | - | - | - |
| SAN JOSE SCALE | 3/19 | - | - | - | - | 2.8 | 250 |

Codling moth degree days are 200 after the 3/26 biofix. UC IPM suggests 300 degree days if you plan to treat the over wintering generation. Our CM trap catches remain low. We caught one male and 3 females on 4/19 and no moths on 4/23.

Additional Pest management information is available at the UC IPM website.

<http://www.ipm.ucdavis.edu>.

Richard P. Buchner and Cyndi K. Gilles
 Orchard Advisor & Research Associate
 UC Cooperative Extension Tehama County
 (530) 527-3101