

University of California IPFP, IPM, and WPMA Pest Update #12
Tehama County 6/4/07

IPFP = Integrated Prune Farming Practices
 IPM = Integrated Pest Management
 WPMA = Walnut Pest Management Alliance

INSECT	FIRST BIOFIX	SECOND BIOFIX	THIRD BIOFIX	FOURTH BIOFIX	ACTIVITY MOTHS/DAY	AVG. DEG. DAYS/DAY	DAY DEGREES FROM BIOFIX
CODLING MOTH	3/26	-	-	-	0	19.8	895
ORIENTAL FRUIT MOTH	3/5	5/10	-	-	7.3	24.5	580
PEACH TWIG BORER	4/2	-	-	-	4.2	19.8	848
NAVEL ORANGE WORM	4/30	-	-	-	0	15.0	437
SAN JOSE SCALE	3/19	-	-	-	-	18.8	907

Codling moth degree days are 895 after the 3/26 biofix. In the walnuts that we monitor we have not measured significant moth increases typical for the “1B” Trap increases in the 900 to 1000 DD range should be the next generation.

We typically catch our first Walnut Husk Flies about July first in the Dairyville area. It’s not too early to order traps and get prepared for Walnut Husk Fly monitoring. We plan to get traps up by mid June.

Additional Pest management information is available at the UC IPM website.
<http://www.ipm.ucdavis.edu>.

Richard P. Buchner and Cyndi K. Gilles
 Orchard Advisor & Research Associate
 UC Cooperative Extension Tehama County
 (530) 527-3101

