

University of California IPFP, IPM, Pest Update #5
Tehama County 4/21/08

IPFP = Integrated Prune Farming Practices
IPM = Integrated Pest Management

INSECT	FIRST BIOFIX	SECOND BIOFIX	THIRD BIOFIX	FOURTH BIOFIX	ACTIVITY MOTHS/DAY	AVG. DEG. DAYS/DAY	DAY DEGREES FROM BIOFIX
CODLING MOTH	3/24	-	-	-	6.8	7.0	195
ORIENTAL FRUIT MOTH	3/5	-	-	-	0	10.3	455
PEACH TWIG BORER	4/10	-	-	-	.5	7.0	112
NAVEL ORANGE WORM	-	-	-	-	-	-	-
SAN JOSE SCALE	-	-	-	-	-	-	-

In the Cone Grove area walnut orchard that we have historically monitored we have a clear biofix for Codling moth on 3/24/08 and good sustained catches through 4/21. We caught our first female 4/14. We are currently at 195 Degree Days Following Biofix. Conventional insecticide applications are usually timed at 300 Degree Days. If using an Insect Growth Regulator Like Intrepid, application is suggested at 200 DD. Detailed Information on Codling moth spray timing is available at the UC IPM website listed below.

Additional Pest management information is available at the UC IPM website.
<http://www.ipm.ucdavis.edu>.

Richard P. Buchner and Cyndi K. Gilles
Orchard Advisor & Research Associate
UC Cooperative Extension Tehama County
(530) 527-3101