

University of California IPFP, IPM, Pest Update #16
Tehama County 7/7/08

IPFP = Integrated Prune Farming Practices
IPM = Integrated Pest Management

INSECT	FIRST BIOFIX	SECOND BIOFIX	THIRD BIOFIX	FOURTH BIOFIX	ACTIVITY MOTHS/DAY	AVG. DEG. DAYS/DAY	DAY DEGREES FROM BIOFIX
CODLING MOTH	3/24	6/23	-	-	1.5	26.5	348
ORIENTAL FRUIT MOTH	3/5	5/19	-	-	0	31.8	1282
PEACH TWIG BORER	4/10	6/16	-	-	8.6	26.5	520
NAVEL ORANGE WORM	-	-	-	-	0	-	-
SAN JOSE SCALE	4/25	-	-	-	-	25.8	1398

Codling moth in the orchard we use for reporting are at 348 Degree days. The second Biofix was set on 6/23. In the 3 walnut orchards we monitor, we have seen sustained trap increases beginning between 6/23 and 6/30.

It's not too early to start thinking about getting Walnut Husk Fly traps up. We caught our first Husk Fly (male) on 6/13 in the Dairyville area and caught an additional male on 7/3. Remember that the females have blond front legs.

Additional Pest management information for Codling moth and Husk fly is available at the UC IPM website. <http://www.ipm.ucdavis.edu>.

Richard P. Buchner and Cyndi K. Gilles
Orchard Advisor & Research Associate
UC Cooperative Extension Tehama County
(530) 527-3101