

University of California IPFP, IPM, Pest Update #28  
Tehama County 9/29/08

IPFP = Integrated Prune Farming Practices  
IPM = Integrated Pest Management

| INSECT              | FIRST BIOFIX | SECOND BIOFIX | THIRD BIOFIX | FOURTH BIOFIX | ACTIVITY MOTHS/DAY | AVG. DEG. DAYS/DAY | DAY DEGREES FROM BIOFIX |
|---------------------|--------------|---------------|--------------|---------------|--------------------|--------------------|-------------------------|
| CODLING MOTH        | 3/24         | 6/23          | 8/7          | -             | 0                  | 20.6               | 1245                    |
| ORIENTAL FRUIT MOTH | 3/5          | 5/19          | 6/26         | -             | 0                  | 25.6               | 2820                    |
| PEACH TWIG BORER    | 4/10         | 6/16          | 7/28         | -             | 6.0                | 20.6               | 1489                    |
| NAVEL ORANGE WORM   | -            | -             | -            | -             | 0                  | -                  | -                       |
| SAN JOSE SCALE      | 4/25         | -             | -            | -             | -                  | 1938               | 3382                    |

Additional Pest management information for Codling moth and Husk fly is available at the UC IPM website. <http://www.ipm.ucdavis.edu>.

Richard P. Buchner and Cyndi K. Gilles  
Orchard Advisor & Research Associate  
UC Cooperative Extension Tehama County  
(530) 527-3101