

University of California IPFP, IPM, Pest Update #12
Tehama County 6/15/09

IPFP = Integrated Prune Farming Practices
IPM = Integrated Pest Management

INSECT	FIRST BIOFIX	SECOND BIOFIX	THIRD BIOFIX	FOURTH BIOFIX	ACTIVITY MOTHS/DAY	AVG. DEG. DAYS/DAY	DAY DEGREES FROM BIOFIX
CODLING MOTH	3/30	-	-	-	0	18.8	1220
ORIENTAL FRUIT MOTH	3/16	5/18	-	-	.38	23.8	727
PEACH TWIG BORER	4/20	-	-	-	.5	18.8	1017
NAVEL ORANGE WORM	-	-	-	-	0	-	-
SAN JOSE SCALE	-	-	-	-	0	-	-

Additional Pest management information for Codling moth is available at the UC IPM website <http://www.ipm.ucdavis.edu>. The Second flight is predicted to start as early as 800 DD and as late as 1300 DD. The Average is about 1000 DD. We haven't seen a large trap catch increase but slight activity on 6/11 suggesting that may be the second biofix. We will confirm sustained trap catches on Thursday.

It's not too early to start thinking about getting Walnut Husk Fly traps ordered and into orchards. We usually catch flies about the first week in July. We have caught no WHF yet.

Richard P. Buchner and Cyndi K. Gilles
Orchard Advisor & Research Associate
UC Cooperative Extension Tehama County
(530) 527-3101