

University of California IPFP, IPM, Pest Update #13
Tehama County 6/22/09

IPFP = Integrated Prune Farming Practices
IPM = Integrated Pest Management

INSECT	FIRST BIOFIX	SECOND BIOFIX	THIRD BIOFIX	FOURTH BIOFIX	ACTIVITY MOTHS/DAY	AVG. DEG. DAYS/DAY	DAY DEGREES FROM BIOFIX
CODLING MOTH	3/30	6/11	-	-	6.00	24.0	239
ORIENTAL FRUIT MOTH	3/16	5/18	-	-	.56	29.5	928
PEACH TWIG BORER	4/20	6/18	-	-	1.2	24.0	96
NAVEL ORANGE WORM	-	-	-	-	0	-	-
SAN JOSE SCALE	-	-	-	-	0	-	-

Additional Pest management information for Codling moth is available at the UC IPM website <http://www.ipm.ucdavis.edu>. We monitor three walnut orchards for Codling Moth. Until today, trap catches have been relatively low making it difficult to pick a clear second biofix. Looking at all our data we selected the second biofix for 6/11 at 1145 degree days. Catches have been low until today when we caught 24 moths with a combo lure and 21 moths with a pheromone lure over a 4 day period, suggesting significant moth activity has started.

It's not too early to start thinking about getting Walnut Husk Fly traps into orchards. We caught our first male walnut Husk Fly today 6/22/09.

Richard P. Buchner and Cyndi K. Gilles
Orchard Advisor & Research Associate
UC Cooperative Extension Tehama County
(530) 527-3101