

University of California IPFP, IPM, Pest Update #21
Tehama County 8/17/09

IPFP = Integrated Prune Farming Practices
IPM = Integrated Pest Management

INSECT	FIRST BIOFIX	SECOND BIOFIX	THIRD BIOFIX	FOURTH BIOFIX	ACTIVITY MOTHS/DAY	AVG. DEG. DAYS/DAY	DAY DEGREES FROM BIOFIX
CODLING MOTH	3/30	6/11	7/23	-	1.5	26.5	658
ORIENTAL FRUIT MOTH	3/16	5/18	6/25	7/27	.38	31.8	654
PEACH TWIG BORER	4/20	6/18	7/31	-	3.4	26.5	431
NAVEL ORANGE WORM	-	-	-	-	0	-	-
SAN JOSE SCALE	-	-	-	-	0	-	-

We had a third Codling moth biofix on 7/23 and are currently at 658 day degrees. Additional Pest management information for Codling moth is available at the UC IPM website <http://www.ipm.ucdavis.edu>.

Watch walnut husk fly traps carefully and treat when you find first females with eggs. Females have blond front legs and the eggs look like grains of rice when you squish the abdomen. We saw large trap catches last week and treated the orchards we monitor. Due to the sprays, Husk fly counts are down this week. WHF may be down but they are not out so watch traps carefully for any new emergence.

Richard P. Buchner and Cyndi K. Gilles
Orchard Advisor & Research Associate
UC Cooperative Extension Tehama County
(530) 527-3101